

Sub A1

- Sub A2

8. A method according to claim 1, 3-7 whereby said dendritic cells are infected with one or more recombinant viruses encoding the antigen(s) of interest before activating said dendritic cells in the presence of a glucocorticoid hormone.
9. A method according to claim 1, 3-8 whereby said dendritic cells are incubated with one or more recombinant proteins or large (> 20 amino acids) synthetic peptides representing the antigen(s) of interest before activating said dendritic cells in the presence of a glucocorticoid hormone.
10. A method according to claim 1, 3 or 9 whereby said dendritic cells are incubated with cells or cell homogenate containing the antigen(s) of interest before activating said dendritic cells in the presence of a glucocorticoid hormone.
11. A method according to claim 1, 3-10 whereby said dendritic cells are loaded with synthetic peptides representing the antigen(s) of interest after activating said dendritic cells in the presence of a glucocorticoid hormone.
12. A method according to claim 1, 3-11 whereby said dendritic cells, after activation in the presence of a glucocorticoid hormone, secrete interleukin-10.
13. A method for obtaining a dendritic cell capable of tolerising a T-cell for an antigen comprising providing said dendritic cell with a glucocorticoid hormone, activating said dendritic cell and providing said dendritic cell with said antigen.
14. A method according to anyone of claims 1, 3-13, wherein said dendritic cell and/or a precursor thereof is provided with said glucocorticoid hormone in vitro.
15. A method according to anyone of claims 1, 3-14, wherein said T-cell is a T-helper cell.
16. An isolated dendritic cell prepared according to anyone of claims 1, 3-15 capable of functionally modifying an antigen-specific T-cell with respect to the response to said antigen.

001260 02499560

Sub A2

5

5

1

10

15.

15.

20

25

30

30

35

add A5  
82

00027